

Metronidazole 5 mg/ml solution for infusion

Metronidazole

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Metronidazole is and what it is used for
2. What you need to know before you take Metronidazole
3. How to take Metronidazole
4. Possible side effects
5. How to store Metronidazole
6. Contents of the pack and other information

1. WHAT METRONIDAZOLE IS AND WHAT IT IS USED FOR

Metronidazole belongs to a group of medicines known as antibiotics and is used to treat severe infections caused by bacteria that can be killed by the active substance metronidazole.

You may be given Metronidazole for the treatment of any of the following diseases:

- Infections of the blood, brain, lung, bones, genital tract, pelvic area and stomach

If required, your treatment may be supplemented by other antibiotics.

Metronidazole may be given as a preventive measure prior to operations associated with a higher risk of infection with what are known as anaerobic bacteria, mainly in gynaecology or surgery on stomach and gut.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE METRONIDAZOLE

Do not take Metronidazole:

if you are allergic (hypersensitive) to metronidazole, other similar substances or any of the other ingredients of Metronidazole (listed in Section 6).

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Metronidazole if you have:

- severe liver damage,
- a blood formation disorder or
- a disease of brain, spinal cord or nerves

Therefore, your doctor will very carefully determine whether you should be treated with Metronidazole.

If convulsive fits or any other nerve affections (e.g. numbness in limbs) become apparent during therapy, your treatment will promptly be revised. Treatment must be stopped or revised immediately if you get severe diarrhoea which may be due to a severe large bowel disease called "pseudomembranous colitis" (see also section 4.)

As prolonged use of metronidazole may impair blood formation (see section "Possible side effects"), your blood counts will be monitored during treatment.

If you received this medicine your urine may be darkened.

Cases of severe liver toxicity/acute liver failure, including cases with a fatal outcome, in patients with Cockayne syndrome have been reported with product containing metronidazole.

If you are affected by Cockayne syndrome, your doctor should also monitor your liver function frequently while you are being treated with metronidazole and afterwards.

Tell your doctor immediately and stop taking metronidazole if you develop:

- Stomach pain, anorexia, nausea, vomiting, fever, malaise, fatigue, jaundice, dark urine, putty or mastic coloured stools or itching.

Treatment with Metronidazole should not usually be continued for longer than 10 days; the treatment period will only be extended in exceptional circumstances and if absolutely necessary. Repeat therapy with metronidazole will be restricted to cases where this is absolutely necessary. In such a case, you will be monitored particularly carefully.

Other medicines and Metronidazole

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

Amiodarone (used to treat irregular heartbeat)

When you receive this medicine, your heart function should be monitored. You should see your doctor if you notice any heart function abnormalities, dizziness or fainting.

Barbiturates (the active substance in sleeping pills)

The duration of action of metronidazole is reduced by phenobarbital; your metronidazole dose may therefore have to be increased.

Busulfan

Metronidazole should not be given to patients receiving busulfan because in that case toxic effects are more likely to occur.

Carbamazepine (a drug for the treatment of epilepsy)

This combination also warrants caution because metronidazole may increase the duration of action of carbamazepine.

Cimetidine (a drug for the treatment of stomach disorders)

Cimetidine may reduce the elimination of metronidazole in isolated cases and subsequently leads to increased serum metronidazole concentrations.

Coumarin derivatives (drugs that inhibit blood clotting)

Metronidazole may enhance the blood clotting inhibition brought about by coumarins. So if you are taking a medicine that inhibits blood clotting (for example warfarin), you may need less of it during treatment with metronidazole.

Cyclosporin (a drug used to suppress undesirable immune responses)

When cyclosporin is given together with metronidazole, the blood levels of cyclosporin may increase; your doctor will therefore have to adjust your cyclosporin dose as appropriate.



Metronidazole 5 mg/ml solution for infusion

Metronidazole

The following information is intended for healthcare professionals only:

Posology and method of administration

The dosage is adjusted according to the patient's individual response to therapy, her/his age and body weight and according to nature and severity of the disease.

The following dosage guidelines should be followed:

Adults and adolescents:

Treatment of anaerobic infections
500 mg (100 ml) every 8 hours. Alternatively 1000 mg – 1500 mg may be given daily as a single dose.

The duration of therapy is dependent on the effect of the treatment. In most cases a treatment course of 7 days will be sufficient. If clinically indicated, treatment may be continued beyond this time although a duration of 10 days should not normally be exceeded. (See also section 4.4.)

Disulfiram (used in alcohol withdrawal therapy)

If you are taking disulfiram, you must not be given metronidazole, or disulfiram must be stopped. Combined use of these two drugs may lead to states of confusion up to the point of a serious mental disorder (psychosis).

Drugs containing alcohol

Please refer to section 'Using Metronidazole' with food and drink.

Fluorouracil (an anticancer drug)

The daily dose of Fluorouracil may have to be reduced when giving it together with Metronidazole because metronidazole may lead to an increase of the blood level of Fluorouracil.

Lithium (used to treat mental illness)

Treatment with lithium preparations requires particularly careful monitoring during treatment with metronidazole, and the dose of the lithium preparation may need to be re-adjusted. Lithium treatment should be tapered or withdrawn before administration of metronidazole.

Mycophenolate mofetil (used for the prevention of rejection reactions after organ transplant)

Its effect may be weakened by metronidazole, so careful monitoring of the effect of the medicine is recommended.

Phenytoin (a drug for the treatment of epilepsy)

If you are taking phenytoin, your doctor will treat you with metronidazole only with caution because metronidazole may increase the duration of action of phenytoin. On the other hand, phenytoin may reduce the effect of metronidazole.

Tacrolimus (used to suppress unwanted immune reactions)

The blood levels of this agent and your kidney function should be checked when starting and stopping treatment with metronidazole.

Metronidazole with food, drink and alcohol

Alcohol

You must not drink any alcoholic beverages or drugs containing alcohol while you are being given metronidazole and up to 48 hours afterwards because this may cause intolerance reactions such as dizziness and vomiting.

Pregnancy breast-feeding and fertility

Fertility

Animal studies only indicate a potential negative influence of metronidazole on the male reproductive system if high doses lying well above the maximum recommended dose for humans were administered.

Pregnancy

If you are pregnant, your doctor will not treat you with metronidazole unless she/he considers this absolutely necessary.

Breast-feeding

You should not breast-feed during treatment with metronidazole and not resume nursing for another 2–3 days thereafter because metronidazole passes into breast milk.

Driving and using machines

While taking Metronidazole you may feel sleepy, dizzy, confused, see or hear things that are not there (hallucinations), have fits (convulsions) or temporary eyesight problems (such as blurred or double vision). If this happens, do not drive or use any machinery or tools.

Metronidazole contains sodium

This medicine contains 14 mmol (or 322 mg) sodium per 100 ml. To be taken into consideration by patients on a controlled sodium diet

3. HOW TO TAKE METRONIDAZOLE

Dosage

Dosage depends on the nature and severity of your illness, your age and body weight, and your individual response to treatment.

The following dosages are usually prescribed:

Adults and adolescents

Treatment of infections:

Adults

You will be given 100 ml of the medicine (500 mg of metronidazole) every 8 hours.

In most cases treatment will take 7 days. Only exceptionally treatment may be continued beyond this time, although a duration of 10 days should not normally be exceeded.

The dose will be the same for patients with kidney diseases.

For patients with liver diseases, lower doses may be required.

If you were treated by artificial kidney your doctor will schedule your infusion after dialysis has been finished. No routine dose adjustment is necessary.

Prevention of infections that might occur after operations

When used for prevention of infection in surgery, you may be given 500 mg of the medicine before the operation. The dose will be repeated 8 and 16 hours after the operation.

The Elderly:

Your doctor will give you this medicine only with special caution.

Children

Dosing in children is based on body weight (BW).

Prophylaxis against post-operative infection caused by anaerobic bacteria: 500 mg, with administration completed approximately one hour before surgery. The dose is repeated after 8 and 16 hours.

The Elderly:

Caution is advised in the elderly, particularly at high doses, although there is limited information available on modification of dosage.

Paediatric population

Treatment of anaerobic infections

Children > 8 weeks to 12 years of age:
The usual daily dose is 20 – 30 mg per kg BW per day as a single dose or divided into 7.5 mg per kg BW every 8 hours. The daily dose may be increased to 40 mg per kg BW, depending on the severity of the infection.

Treatment of infections:

Age	Dosage
8 weeks to 12 years	20 – 30 mg of metronidazole per kg BW per day as a single dose or divided into 7.5 mg of metronidazole per kg BW every 8 hours. The daily dose may be increased to 40 mg of metronidazole per kg BW if infection is severe.
Under 8 weeks	15 mg of metronidazole per kg BW as a single dose daily or divided into 7.5 mg per kg BW every 12 hours.
Newborns of less than 40 weeks gestational age	As metronidazole may accumulate in these patients during the first week of life, the concentration of metronidazole in the blood will be checked after a few days of treatment.

Usually treatment will take 7 days.

Prevention of infections that might occur after operations:

Age	Dosage
Less than 12 years	20 – 30 mg of metronidazole per kg BW as a single dose given 1 – 2 hours before surgery
Newborns of less than 40 weeks gestational age	10 mg of metronidazole per kg BW as a single dose before surgery

Method of administration and duration of treatment

Metronidazole is administered through a drip directly into a vein (intravenous infusion)

The infusion of one bottle usually takes 60 minutes, but it should not be done within less than 20 minutes.

The entire metronidazole treatment period is usually 7 days and must not exceed 10 days unless this is absolutely necessary (see also “Take special care with Metronidazole”).

If you are concurrently receiving other antibiotics your doctor will give you those medicines separately

If you take more Metronidazole than you should

Undesirable effects, as described in the next section, may occur as signs or symptoms of an overdose. Single oral doses of metronidazole, up to 12 g have been reported in suicide attempts and accidental overdoses. Symptoms were limited to vomiting, ataxia and slight disorientation.

There is no known specific antidote or specific treatment of a massive overdose, but metronidazole can be removed by dialysis (that is treatment with artificial kidney) from the body.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Metronidazole can cause side effects, although not everybody gets them.

Side effects occur mostly at high doses or with prolonged use.

Talk to your doctor straight away if you notice any of the following side effects:

Rare (may affect up to 1 in 1,000 people):

- Severe persistent diarrhoea (possibly a symptom of a severe bowel infection called pseudomembranous colitis, see below)
- Severe acute hypersensitivity reactions up to allergic shock

Very rare (may affect up to 1 in 10,000 people):

- White blood cell and platelet counts may decrease during treatment (granulocytopenia, agranulocytosis, pancytopenia, thrombocytopenia)
- Hepatitis (liver inflammation), jaundice, inflammation of the pancreas
- Brain disorders, lack of coordination
- Brain fever not caused by bacteria (aseptic meningitis)
- Severe inflammatory rash on mucous membranes and the skin with fever, redness and blistering, in extremely rare cases up to skin detachment over extended areas (Stevens-Johnson Syndrome)

Not known (frequency cannot be estimated from the available data):

- Mild to moderate hypersensitivity reactions, swelling of your face, mouth, throat and/or tongue (angioedema)
- Gaze spasm, damage or inflammation of the nerves of your eyes
- Reduced white blood cell, count (leucopenia), severe anaemia (aplastic anaemia)
- Seizures, nervous disorders such as numbness, pain, furry sensation or tingling in the arms or legs
- Toxic epidermal necrolysis

Other side effects include

Common (may affect up to 1 in 10 people)

- Infections with yeasts (e.g. genital infections)

Uncommon (may affect up to 1 in 100 people)

- Darkened urine (due to a metabolite of metronidazole)

Rare (may affect up to 1 in 1,000 people)

- Changes in ECG

Very rare (may affect up to 1 in 10,000 people):

- Psychotic disorders, including states of confusion, hallucination
- Headache, dizziness, drowsiness, fever, disturbance of sight and movement, giddiness, speech defects, convulsions
- Visual disturbances, e.g. double vision, short-sightedness
- Liver function disorders (such as elevated serum levels of certain enzymes and bilirubin)
- Allergic skin reactions like itching, hives
- Joint and muscle pain

Not known (frequency cannot be estimated from the available data):

- Sickness, feeling sick, diarrhoea, inflammation of tongue or mouth, belching and bitter taste, metallic taste, pressure above the stomach, furry tongue
- Difficulty swallowing
- Anorexia
- Sad (depressed) mood
- Sleepiness or sleeplessness, muscle twitching
- Reddening and itching of the skin (erythema multiforme)
- Vein wall irritation (to the point of inflamed veins and thrombosis) after intravenous administration, states of weakness, fever

Emergency management of pseudomembranous enterocolitis.

In the event of severe persistent diarrhoea, you must promptly inform your doctor because this may be due to pseudomembranous colitis, a serious condition that must be treated immediately. Your doctor will stop metronidazole and provide appropriate treatment.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse.

This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via the Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE METRONIDAZOLE

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bag and the carton after EXP. The expiry date refers to the last day of that month.

Do not store above 25 °C.

Do not refrigerate.

Keep the bag in the original packaging (aluminium overwrap) in order to protect from light.

After first opening the medicinal product should be used immediately

Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Metronidazole contains

The active substance is metronidazole.

Each bag of 100 ml of solution for infusion contains 500 mg metronidazole.

Each ml of solution for infusion contains 5 mg metronidazole.

Other ingredients are: Sodium chloride, disodium phosphate, anhydrous, citric acid, anhydrous and water for injections.

What Metronidazole looks like and contents of the pack

Metronidazole is a clear, colourless or slightly yellowish aqueous solution infusion

Metronidazole is contained in polypropylene bags closed with a twist off port, inside an aluminum overwrap, containing 100 ml of solution.

This medicine is marketed as individual packs or packs of 10 bags.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Hikma Farmacêutica (Portugal) S.A.

Estrada do Rio da Mó, nº8, 8A e 8B

Fervença

2705-906 Terrugem SNT

Portugal

Tel.:+ 351 219 608 410

Email: portugalgeral@hikma.com

Distributed by:

Consilient Health (UK) Ltd.

No.1 Church Road,

Richmond upon Thames,

Surrey. TW9 2QE

This leaflet was last revised in April 2020.



Neonates and infants < 8 weeks of age:

15 mg per kg BW as a single dose daily or divided into 7.5 mg per kg BW every 12 hours.

In newborns with a gestational age < 40 weeks, accumulation of metronidazole can occur during the first week of life; therefore the concentrations of metronidazole in serum should preferably be monitored after a few days therapy.

Duration of treatment is usually 7 days.

Prophylaxis against postoperative infections caused by anaerobic bacteria:

Children < 12 years:

20 – 30 mg/kg BW as a single dose given 1 – 2 hours before surgery

Newborns with a gestational age < 40 weeks:

10 mg/kg BW as a single dose before surgery

Patients with renal insufficiency

Limited data are available in this population. These data do not indicate the need for dose reduction (see section 5.2.)

In patients undergoing haemodialysis the conventional dose of metronidazole should be scheduled after haemodialysis on dialysis days to compensate the removal of metronidazole during the procedure.

No routine dose adjustment is necessary in patients with renal failure undergoing intermittent peritoneal dialysis (IDP) or continuous ambulatory peritoneal dialysis (CAPD).

Patients with hepatic insufficiency

As serum half-life is prolonged and plasma clearance is delayed in severe hepatic insufficiency, patients with severe liver disease will require lower doses (see section 5.2).

In patients with hepatic encephalopathy, the daily dosage should be reduced to one third and may be administered once daily (see section 4.4).

Method of administration

Intravenous use.

The contents of one bottle are to be infused slowly i.v., i.e. 100 ml max. over not less than 20 minutes, but normally over one hour.

Concurrently prescribed antibiotics are to be administered separately.